

## Somatostatin rabbit pAb

Cat No.: ES3479

For research use only

## Overview

Product Name Somatostatin rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human Somatostatin. AA

range:10-59

**Specificity** Somatostatin Polyclonal Antibody detects

endogenous levels of Somatostatin protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Store at -20°C.** Avoid repeated freeze-thaw cycles.

Protein Name Somatostatin

Gene Name SST

Cellular localization Secreted.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

ClonalityPolyclonalConcentration1 mg/mlObserved band17kDHuman Gene ID6750Human Swiss-Prot NumberP61278

**Alternative Names** SST; Somatostatin; Growth hormone

release-inhibiting factor

Background The hormone somatostatin has active 14 aa and 28


aa forms that are produced by alternate cleavage of the single preproprotein encoded by this gene. Somatostatin is expressed throughout the body and

inhibits the release of numerous secondary






hormones by binding to high-affinity G-protein-coupled somatostatin receptors. This hormone is an important regulator of the endocrine system through its interactions with pituitary growth hormone, thyroid stimulating hormone, and most hormones of the gastrointestinal tract. Somatostatin also affects rates of neurotransmission in the central nervous system and proliferation of both normal and tumorigenic cells. [provided by RefSeq, Jul 2008],




Western blot analysis of lysates from 1) PC-12, 2) A549 ,3) Hela cells, (Green) primary antibody was diluted at 1:1000, 4°over night, secondary antibody(cat:RS23920)was diluted at 1:10000, 37° 1hour. (Red.) Actin  $\beta$  Monoclonal Antibody(5B7) (cat:YM3028) ant



Western Blot analysis of various cells using Somatostatin Polyclonal Antibody







Western Blot analysis of A549 cells using Somatostatin Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human Gastric fundus. 1, Antibody was diluted at 1:200(4° overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

