ELK Biotechnology

Recombinant Human NPR3 (C-Fc)

Catalog #	EPT167
Expression Host	Human Cells
DESCRIPTION	Recombinant Human Atrial Natriuretic Peptide Receptor 3 is produced by our Mammalian expression system and the target gene encoding Thr24-Glu481 is expressed with a Fc tag at the C-terminus.
Accession	P17342
Synonyms	ANP-C; ANPR-C; NPR3; NPRC; NPR-C;ANPRC; C5orf23
Mol Mass	77.5 KDa
AP Mol Mass	90-100 KDa, reducing conditions
Purity	Greater than 95% as determined by reducing SDS-PAGE.
Endotoxin	Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
FORMULATION	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
RECONSTITUTION	Always centrifuge tubes before opening.Do not mix by vortex or pipetting.

+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road,Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C

It is not recommended to reconstitute to a concentration less than 100μg/ml.

Dissolve the lyophilized protein in distilled water.

Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SHIPPING

The product is shipped at ambient temperature.

Upon receipt, store it immediately at the temperature listed below.

STORAGE

Lyophilized protein should be stored at < -20 ° C, though stable at room temperature for 3 weeks.

Reconstituted protein solution can be stored at 4-7°C for 2-7 days.

Aliquots of reconstituted samples are stable at < -20° C for 3 months.

BACKGROUND

Atrial Natriuretic Peptide Receptor-3 (NPR3), also known as NPRC or ANPR-C, is one of the three natriuretic peptide receptors, is a type I transmembrane glycoprotein. The natriuretic system is key to the maintenance of vascular tone and cardiovascular homeostasis. Receptor for the natriuretic peptide hormones, binding with similar affinities atrial natriuretic peptide NPPA/ANP, brain

natriuretic peptide NPPB/BNP, and C-type natriuretic peptide NPPC/CNP. May function as a clearance receptor for NPPA, NPPB and NPPC, regulating their local concentrations and effects. Osteocrin was found to be a specific ligand to NPR3. NPR3 is necessary for Osteocrin to regulate femoral, tibial, and metatarsal bone elongation.

SDS-PAGE